

# IOWA STATE UNIVERSITY

## Digital Repository

---

Chemistry Publications

Chemistry

---

6-1-2019

## Efficient, Scalable Syntheses of Ginkgolic Acids

Joshua L. Alterman

*Iowa State University*, [alterman@iastate.edu](mailto:alterman@iastate.edu)

George A. Kraus

*Iowa State University*, [gakraus@iastate.edu](mailto:gakraus@iastate.edu)

Follow this and additional works at: [https://lib.dr.iastate.edu/chem\\_pubs](https://lib.dr.iastate.edu/chem_pubs)

 Part of the [Organic Chemistry Commons](#)

The complete bibliographic information for this item can be found at [https://lib.dr.iastate.edu/chem\\_pubs/1166](https://lib.dr.iastate.edu/chem_pubs/1166). For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

---

This Article is brought to you for free and open access by the Chemistry at Iowa State University Digital Repository. It has been accepted for inclusion in Chemistry Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

## Efficient, Scalable Syntheses of Ginkgolic Acids

### Abstract

Ginkgolic acids were directly synthesized from 6-methylsalicylic acid using a sequence involving protection of the phenol and acid as methoxymethyl ethers and esters, lateral alkylation and deprotection under mild aqueous acid conditions.

### Keywords

lateral alkylation, anti-HIV, direct synthesis

### Disciplines

Organic Chemistry

### Comments

This article is published as Alterman, Joshua L., and George A. Kraus. "Efficient, Scalable Syntheses of Ginkgolic Acids." *Natural Product Communications* 14, no. 6 (2019). DOI: [10.1177%2F1934578X19851348](https://doi.org/10.1177%2F1934578X19851348). Posted with permission.

### Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial 4.0 License](https://creativecommons.org/licenses/by-nc/4.0/)

# Efficient, Scalable Syntheses of Ginkgolic Acids

Joshua L. Alterman<sup>1</sup> and George A. Kraus<sup>1</sup>

Natural Product Communications  
June 2019: 1–3  
© The Author(s) 2019  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/1934578X19851348  
journals.sagepub.com/home/npx



## Abstract

Ginkgolic acids were directly synthesized from 6-methylsalicylic acid using a sequence involving protection of the phenol and acid as methoxymethyl ethers and esters, lateral alkylation and deprotection under mild aqueous acid conditions.

## Keywords

lateral alkylation, anti-HIV, direct synthesis

Received: January 24th, 2019; Accepted: February 21st, 2019.

Ginkgolic acid C15:1 (**1**) and ginkgolic acid C15:0 (**2**) are natural acids isolated from *Ginkgo biloba* (Figure 1).<sup>1</sup> Ginkgolic acid is a component of a botanical extract which shows pleiotropic effects including antitumor and anti-HIV activities.<sup>2</sup> The synthesis of (**1**) has been reported by Martin in 2018 using a Heck-based strategy.<sup>3</sup> Several researchers used Wittig-based strategies.<sup>4–7</sup> The lateral alkylation strategy presented herein, also used by Tyman,<sup>8</sup> utilizes commercially available and stable reagents such as (**3**) and is amenable to scale up.

Protection of 6-methyl salicylic acid (**3**) with in situ derived chloromethyl methyl ether provided ester (**4**) as shown in Scheme 1.<sup>9</sup> A variety of bases were evaluated for the deprotonation of (**4**). Ultimately, treatment of (**4**) with lithium 2,2,6,6-tetramethylpiperidine (LiTMP) in tetrahydrofuran (THF) at  $-78^{\circ}\text{C}$ , followed by the addition of the acetylenic iodide (**5**),<sup>10</sup> and warming to ambient temperature overnight afforded the alkynyl ester (**6**) in 60% yield. Reduction of the alkyne using hydrogen and the Lindlar's catalyst followed by treatment with 1 N HCl to remove the protecting groups afforded (**1**) in 61% yield. The use of the methoxy methyl ether protecting groups increased the solubility of **4** and enables easy global deprotection to **1**.

The reaction of the anion of (**4**) with 1-iodotetradecane followed by deprotection generated (**2**) in 61% yield as shown in Scheme 2. Iodotetradecane was prepared from the commercially available chloride in 90% yield using sodium iodide.

The synthesis of (**1**) and (**2**) in three or four steps constitutes a direct route to these biologically active compounds.<sup>11</sup> This direct pathway will enable the synthesis of (**1**), (**2**), and analogs for biological evaluation.

## Experimental

### Methoxymethyl 2-(Methoxymethoxy)-6-Methylbenzoate (**4**)

To 35 mL DMF was added (**3**) (893 mg, 5.86 mmol) and allowed to stir at  $0^{\circ}\text{C}$  for 15 minutes. Then 5 equivalents of NaH was added in 4 portions. The solution was brought from  $0^{\circ}\text{C}$  to r.t for 30 minutes, then cooled back to  $0^{\circ}\text{C}$ . Chloromethyl methyl ether, (1.4 mL, 3.14 equiv.), was added over 4 additions, every 15 to 20 minutes. Reaction was gently quenched with ice and sat.  $\text{NH}_4\text{Cl}$  and then extracted with ethyl acetate. After drying over  $\text{Na}_2\text{SO}_4$  and concentrating *in vacuo*, the resulting oil was purified using silica gel chromatography (Hexane:Ethyl acetate, 3:7) affording a clear, colorless, oil (**4**) (1.13 g, 80% yield).

$^1\text{H}$  NMR (400 MHz  $\text{CDCl}_3$ ):  $\delta$  (ppm) 7.23 (t,  $J = 8.0$  Hz, 1H), 6.99 (d,  $J = 8.4$  Hz, 1H), 6.87 (d,  $J = 7.6$  Hz, 1H), 5.47 (s, 2H), 5.18 (s, 2H), 3.56 (s, 3H), 3.47 (s, 3H),  $\delta$  2.34 (s, 3H).

### Methoxymethyl-2-(Methoxymethoxy)-6-(Pentadec-8-yn-1-yl) Benzoate (**6**)


To oven dried reaction vessel tetramethylpiperidine (0.26 mL, 1.5 mmol) was added, followed by 6 mL of THF and

<sup>1</sup> Department of Chemistry, Iowa State University, Ames, IA, USA


### Corresponding Author:

George A. Kraus, Department of Chemistry, Iowa State University, Ames, IA 50011, USA.  
Email: gakraus@iastate.edu






**Figure 1.** Structures for ginkgolic acids C15:1 (**1**) and C15:0 (**2**).



**Scheme 1.** Synthesis of **1**.



**Scheme 2.** Synthesis of (**2**).

was cooled to 0°C. Argon was then bubbled through the solution for ~5 minutes. Addition of *n*BuLi (0.60 mL, 1.5 mmol) was done dropwise, and the reaction allowed to reach room temperature (r.t.) for 25 minutes. The solution was then cooled to -78°C where (**4**) (238 mg, 0.991 mmol) in 2 mL of THF was slowly added and reacted for 71 minutes. Then (**5**) (548 mg, 1.7 mmol) in 2 mL of THF was chilled to -78°C, and syringed while cold. The reaction was let warm to room temperature overnight. The reaction was quenched with sat. NH<sub>4</sub>Cl and extracted with ethyl acetate, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated in vacuo. The resulting oil was purified using silica gel chromatography (Hexane:Ethyl acetate, 1:1) affording a clear, orange, oil (**6**) (0.260 g, 60% yield).

<sup>1</sup>H NMR (400 MHz CDCl<sub>3</sub>): δ (ppm) 7.26 (t, *J* = 7.9 Hz, 1H), 7.00 (d, *J* = 8.1 Hz, 1H), 6.89 (d, *J* = 7.7 Hz, 1H), 5.46 (s, 2H), 5.18 (s, 2H), 3.56 (s, 3H), 3.47 (s, 3H), 2.59 (t, *J* = 7.8 Hz, 2H), 2.13 (m, 4H), 1.60 (m, 4H), 1.51-1.22 (m, 16H), 0.88 (t, *J* = 6.5 Hz, 3H).

### (*Z*)-2-Hydroxy-6-(Pentadec-8-en-1-yl)Benzoic acid (**1**)

A flask containing (**6**) (208 mg, 0.48 mmol), in 5 mL of methanol, with 10 μL of quinoline, was sparged with Argon. Lindlar's catalyst was added, and then hydrogen was bubbled through via balloon. Lindlar's catalyst was removed over a pad of Celite, and the filtrate condensed. Treatment of the resulting oil with 1 M HCl in the presence of isopropyl alcohol for 5 hours afforded (**1**) (0.101 g, 61% yield).

<sup>1</sup>H NMR (400 MHz CDCl<sub>3</sub>): δ (ppm) 7.09 (t, *J* = 7.8 Hz, 1H), 6.67 (d, *J* = 8.1 Hz, 1H), 6.62 (d, 1H, *J* = 7.5 Hz), 5.32 (m, 2H), 2.95 (t, *J* = 7.5 Hz, 2H), 1.96 (m, 4H), 1.50 (m, 2H), 1.32-1.18 (m, 17H), 0.84 (t, *J* = 6.6 Hz, 3H).

LRMS (ESI-QTOF) calcd. for C<sub>22</sub>H<sub>34</sub>O<sub>3</sub> [M-H]<sup>-</sup> 345.2435, found 345.2445.

### 2-Hydroxy-6-Pentadecylbenzoic Acid (**2**)

To an oven dried flask tetramethylpiperidine (0.34 mL, 2.0 mmol) was added, followed by 2 mL of THF and was cooled to 0°C. Argon was then bubbled through the solution for ~5 minutes. Addition of *n*BuLi (0.80 mL, 2.0 mmol) was done dropwise, and the flask allowed to reach r.t. for 26 minutes. It was then cooled to -78°C where (**4**) (291 mg, 1.21 mmol) in 2 mL of THF was slowly added and reacted for 70 minutes. 1-Iodotetradecane (778 mg, 2.4 mmol) in 2 mL of THF, was chilled to -78°C, and syringed while cold. The reaction was let warm to room temperature overnight. The reaction was acidified with 1 M HCl in the presence of isopropyl alcohol for 5 hours and extracted with ethyl acetate, dried over Na<sub>2</sub>SO<sub>4</sub>, and then concentrated in vacuo. The resulting solid was purified using silica gel chromatography (Hexane:Ethyl acetate, 1:1) affording a brownish orange solid (**2**) (0.257 g, 61% yield).

<sup>1</sup>H NMR (400 MHz CDCl<sub>3</sub>): δ (ppm) 11.12 (s, OH), 7.29 (t, *J* = 8.2 Hz, 1H), 6.84 (d, *J* = 8.2 Hz, 1H), 6.72 (d, *J* = 7.6 Hz, 1H), 2.88 (t, *J* = 7.5 Hz, 2H), 1.53-1.15 (m, 26H), 0.88 (t, *J* = 7.5 Hz, 3H).

### 1-Iodotetradecane

To an oven dried flask 1-chlorotetradecane (2.35 g, 14.4 mmol) was added, followed by 50 mL of acetone and NaI (3.01 g, 20.1 mmol), and refluxed for 48 hours. Then the reaction was cooled to 0°C and extracted with hexane and concentrated *in vacuo*. Residual yellow discoloration was removed over a pad of silica affording a clear, colorless, oil (4.20 g, 90% yield).

<sup>1</sup>H NMR (400 MHz CDCl<sub>3</sub>): δ (ppm) 3.18 (t, *J* = 7.1 Hz, 2H), 1.82 (pent, *J* = 7.2 Hz, 2H), 1.43-1.21 (m, 22H), 0.88 (t, *J* = 6.8 Hz, 3H).

## Acknowledgments

We thank the Antimicrobial Resistance program at ISU for partial support for J.L.A.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) disclosed no financial support for the research, authorship, and/or publication of this article.

## References

1. Durrani AA, Tyman JHP. Long-chain phenols. Part 14. Synthesis of 6-n-alkylsalicylic acids and 3-n-alkylphenols. *J Chem Soc Perkin Trans 1*. 1979;1:2069-2078.
2. Mango D, Weisz F, Nisticò R. Ginkgolic acid protects against A $\beta$ -induced synaptic dysfunction in the hippocampus. *Front Pharmacol*. 2016;7:401-407.
3. Weigel WK, Dennis TN, Kang AS, Perry JJP, Martin DBC. A Heck-based strategy to generate anacardic acids and related phenolic lipids for isoform-specific bioactivity profiling. *Org Lett*. 2018;20(19):6234-6238.
4. Yamagiwa Y, Ohashi K, Sakamoto Y, Hirakawa S, Kamikawa T, Kubo I. Syntheses of anacardic acids and ginkgoic acid. *Tetrahedron*. 1987;43(15):3387-3394.
5. Satoh M, Takeuchi N, Nishimura T, Ohta T, Tobinaga S. Synthesis of anacardic acids, 6-[8(Z), 11(Z)-pentadecadienyl]salicylic acid and 6-[8(Z), 11(Z), 14-pentadecatrienyl]salicylic acid. *Chem Pharm Bull*. 2001;49(1):18-22.
6. Fürstner A, Seidel G. Shortcut syntheses of naturally occurring 5-alkylresorcinols with DNA-cleaving properties. *J Org Chem*. 1997;62(8):2332-2336.
7. Zehnter R, Gerlach H. Synthesis of anacardic acids. *Liebigs Annalen*. 1995;1995(12):2209-2220.
8. Tyman JHP, Visani N. Synthesis of saturated anacardic acids, and alkenyl and Alkynyl Analogues†. *J Chem Res*. 1997;(S):14-15.
9. Linderman RJ, Jaber M, Griedel BD. A simple and cost effective synthesis of chloromethyl methyl ether. *J Org Chem*. 1994;59(21):6499-6500.
10. Fitzpatrick SM, Gries R, Khaskin G, Peach DAH, Iwanski J, Gries G. Populations of the gall midge *Dasineura oxycoccana* on cranberry and blueberry produce and respond to different sex pheromones. *J Chem Ecol*. 2013;39(1):37-49.
11. Fukuda I, Ito A, Hirai G, et al. Ginkgolic acid inhibits protein sumoylation by blocking formation of the E1-SUMO intermediate. *Chem Biol*. 2009;16(2):133-140.